



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.


We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

SOLUTION BY MARCIA LATHAM, New York City.

Let  $BC = a$ ,  $AC = b$ ,  $DC = c$ ,  $\angle DCE = \angle ECB = x$ ,  $\angle ABC = y$ . Then  $\angle DCA = 90 - 2x$ ,  $\angle EDC = 180 - (2x + y)$ , and  $\angle BEC = 180 - (x + y)$ .



$$\sin y = b/40 \quad \text{and} \quad \cos y = a/40. \quad (1)$$

Since  $EC$  bisects  $\angle BCD$ ,

$$a/c = 15/10 = 3/2. \quad (2)$$

In the triangle,  $DCA$ , by the law of sines,

$$c/15 = \cos y / \cos 2x. \quad (3)$$

Combining (1), (2), and (3),  $2a/45 = a/40 \cos 2x$ , or  $\cos 2x = 9/16$ ,  $x = 1/2 \cos^{-1} 9/16 = 27^\circ 53' 8''$ . In the triangle,  $BCD$ , by law of sines,  $\sin (2x + y) / \sin y = a/c$ . Whence, by (1) and (2)

$$\sin (2x + y) = 3b/80. \quad (4)$$

Also,  $a/25 = \sin (2x + y) / \sin 2x$ ; whence, by (4),  $\sin 2x = 15b/16a$ . But  $\sin^2 2x + \cos^2 2x = 1$ . Then  $(15b/16a)^2 + (9/16)^2 = 1$ ; whence

$$b^2 = 7a^2/9. \quad (5)$$

Now, in the triangle,  $ABC$ ,  $a^2 + b^2 = (40)^2$ ; whence, by (5),  $a^2 + 7a^2/9 = 1,600$ ; whence,  $a = 30$  and from (5),  $b = 10\sqrt{7}$ .

Also solved by A. M. HARDING, POLYCARP HANSEN, C. E. HORNE, R. A. JOHNSON, ELMER LATSHAW, E. W. MARTIN, LOUIS ORDANKSY, A. PELLETIER, J. L. RILEY, H. M. ROESER, L. SMITH, D. L. STAMY, H. TSAI, and L. G. WELD.

**2760 [1919, 124]. Proposed by CHARLES N. SCHMALL, New York City.**

In an arithmetical progression, if  $s_n$  be the sum of the first  $n$  terms,  $s_{2n}$  the sum of the first  $2n$  terms, and  $s_{3n}$  the sum of the first  $3n$  terms of the same series, prove that  $s_{2n} - s_n = \frac{1}{3}s_{3n}$ .

SOLUTION BY EMMA M. GIBSON, Springfield (Mo.) High School.

The sum of  $n$  terms of an arithmetical progression is expressed by the formula

$$s_n = \frac{n(a_1 + a_n)}{2},$$

where  $a_1$  and  $a_n$  are the first and  $n$ th terms, respectively.

Hence,  $s_n = n(a_1 + a_n)/2$ ,  $s_{2n} = 2n(a_1 + a_{2n})/2$ , and  $s_{3n} = 3n(a_1 + a_{3n})/2$  are the sums of the first  $n$  terms, the first  $2n$  terms, and the first  $3n$  terms, respectively. Now  $a_{2n} = a_n + nd$ ,  $a_{3n} = a_{2n} + nd = a_n + 2nd$ ,  $d$  being the common difference.

Then  $s_{2n} = 2n(a_1 + a_n + nd)/2$  and  $s_{3n} = 3n(a_1 + a_n + 2nd)/2$  and

$$s_{2n} - s_n = 2n(a_1 + a_n + nd)/2 - n(a_1 + a_n)/2 = n(a_1 + a_n + 2nd)/2 = s_{3n}/3.$$

Also solved by R. D. BOHANNAN, H. L. BRIDGES, JR., H. N. CARLETON, W. F. CHENEY, JR., P. J. DA CUNHA, H. C. GOSSARD, WILLIAM HERBERG, C. N. MILLS, LOUIS O'SHAUGHNESSEY, H. L. OLSON, A. PELLETIER, J. B. REYNOLDS, I. S. SUN, and ELIJAH SWIFT.

**2768 [1919, 171]. Proposed by PAUL CAPRON, U. S. Naval Academy.**

Given the center, a focus, and a point of a conic, construct geometrically the circle of curvature at the point.